

Republic Act 11524 known as Coconut Farmers and Industry Trust Fund (CFITF) Act

The Coconut Farmers and Industry Trust Fund (CFITF) Act, which created the Coconut Farmers and Industry Development Plan was signed on February 26, 2021. The program seeks to (1) Increase the productivity and income of coconut farmers' (2) Poverty alleviation, education, and social equity; and (3) Rehabilitation and modernization of the coconut industry toward farm productivity.

The CFITF shall be maintained for 50 years and used for the coconut farmers' benefit and the coconut industry's development.





Information were lifted from the Technology Guide produced by:
Philippine Coconut Authority
Davao Research Center
Bago Oshiro, Davao City 8000

Reproduced and repackaged by:
Department of Agriculture
AGRICULTURAL TRAINING INSTITUTE
Regional Training Center 02
San Mateo and Cabagan, Isabela

FOR MORE INFORMATION, PLEASE CONTACT:
Philippine Coconut Authority
Cagayan Extension Office
67 Nat'l Highway, Tuguegarao City
(078) 3967579

Agricultural Training Institute Regional Training Center 02 0917-501-4270 rtc2.dcc@ati.da.gov.ph





COCONUT TECHNOLOGY GUIDE
COMMON
SALT FERTILIZATION
ON COCONUT

NOT FOR SALE OR USED FOR COMMERCIAL PURPOSES

Common Salt (Sodium Chloride)

The use of sodium chloride (NaCl) or common salt (CS) as fertilizer is a practical means of increasing coconut production. Salt is the cheapest and best source of chlorine to increase copra yield.

Chlorine deficiency in coconut is widespread in inland areas. A PCA survey conducted nationwide showed that at least 40 coconut producing provinces are severely chlorine deficient.

Advantages

- Accelerates crop growth and development
- Increases copra weight and number of nuts
- Minimizes leaf spot damage
- Environment-friendly under judicious practice


Application

Salt may be applied in three ways:

- Broadcast (in flat to slightly sloping areas)
- Broadcast followed by fork-in at 2-3 inches depth of soil (preferably when salt is combined with nitrogenous fertilizers)
- **Holing** (for hilly-sloping areas distributed in 8-10 inches with 3-5 inches depth around the base of the tree)



Broadcast



Broadcast



Holing



In broadcast application, the fertilizer is placed uniformly over a weeded area around the base of the palm (1 m radius depending on the age of the palm).

Generally, bearing palms are fertilized annually in areas with almost uniform rainfall distribution (Table 1). In areas with distinct wet and dry seasons with uneven rainfall distribution, and those with sandy soils, fertilizers are best applied every six months.

Split application is done at the pre-bearing stage (one to four years) of palms. This practice helps reduce loss of fertilizer nutrients through leaching and run-off and make fertilizer use more effective.

Recommended CS rates for different ages/stages of growth of coconut palms

Age/Stage of palms	Rate of NaCI/ Tree/ year	
Nursery	50 g	
Field planting	100 g	
Six months after planting	150 g	
One year after planting	500 g	
Two years after planting	750 g	
Three years after planting	1.10 kg	
Four years after planting	1.30 kg	
Five years and above	1.50 kg	

Socio-Economic Analysis

Net profits from coconut farming depend much on the fertilizer cost, yield, and copra price.

The use of CS as fertilizer at a rate of 1-2 kg/tree is estimated to give a net return of PhP34,517.97-PhP63,330.79/ha/year (Table 2).

Annual yield and production cost of CS fertilization on coconut

NaCl Rate (kg/tree)	Average Copra Yield (kg/ha)	Gross Returns (PhP)	Production Cost (PhP)	Net Returns (PhP)
0	850	24,140.00	7,970.05	16,169.95
1.0	2,000	56,800.00	22,282.03	34,517.97
1.5	2,500	71,000.00	13,837.12	57,162.88
2.0	2,750	78,100.00	14,569.21	63,330.79